

A holographic superconductor in an external magnetic field

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

JHEP09(2008)121

(<http://iopscience.iop.org/1126-6708/2008/09/121>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 128.125.8.200

The article was downloaded on 09/03/2012 at 22:40

Please note that [terms and conditions apply](#).

A holographic superconductor in an external magnetic field

Tameem Albash and Clifford V. Johnson

*Department of Physics and Astronomy, University of Southern California,
Los Angeles, CA 90089, U.S.A.*

E-mail: albash@usc.edu, johnson1@usc.edu

ABSTRACT: We study a system of a complex charged scalar coupled to a Reissner-Nordström black hole in 3+1 dimensional anti-de Sitter spacetime, neglecting back-reaction. With suitable boundary conditions, the cases of a neutral and purely electric black hole have been studied in various limits and were shown to yield key elements of superconductivity in the dual 2+1 dimensional field theory, forming a condensate below a critical temperature. By adding magnetic charge to the black hole, we immerse the superconductor into an external magnetic field. We show that a family of condensates can form and we examine their structure. For finite magnetic field, they are localized in one dimension with a profile that is exactly solvable, since it maps to the quantum harmonic oscillator. As the magnetic field increases, the condensate shrinks in size, which is reminiscent of the Meissner effect.

KEYWORDS: Gauge-gravity correspondence, AdS-CFT Correspondence.

Contents

1. Introduction	1
2. The background	2
3. The scalar field	3
3.1 Review	3
3.2 Perturbative limit	4
4. Numerical method and results	8
5. Discussion	9
A. Holographic dictionary	11

1. Introduction

Since the early days of AdS/CFT [1–3] (see ref. [4] for a review) it has been tempting to consider applying holographic duality to the study of important strongly coupled phenomena in condensed matter systems, and superconductivity has been high on the list.¹ Since finite temperature in these duals typically implies the presence of a black hole, and since superconductivity requires a condensate to form below a certain critical temperature, the existence of a holographically dual background would seem to require a circumvention of various statements of no-hair theorems (which go back to Wheeler [8] — for a review, see ref. [9]). Generically, the black holes would need to have some kind of scalar hair in order to be dual to a superconductor (the scalar’s asymptotic value would be the condensate vacuum expectation value (vev)).

In a series of studies [10–12], Gubser has presented a case for just the right kind of no-scalar-hair theorem evasion to allow for a superconductor’s dual to exist. The statement seems to be that there do exist solutions that allow for a condensing scalar to be coupled to the black hole if the charge on the black hole is large enough. The scalar couples to (at least) a U(1) under which the black hole is charged, and its condensation breaks the gauge symmetry spontaneously, giving a mass to the gauge field. In particular, if the effective mass in the bulk of the scalar is negative enough, the scalar field develops a non-trivial vev at the boundary, giving the gauge field a non-zero mass.

¹The fact that certain black holes and branes are known to exhibit a sort of Meissner effect [5–7] at zero temperature has always added to the motivation, although it has not been clear how exactly this could be connected to a dual superconductivity.

Studying such solutions is hard to do, since the full equations are coupled and non-linear, and so numerical methods, and a number of limits, have been employed in order to extract the key physics. Gubser has studied [12] the case of non-Abelian Reissner-Nordström black holes condensing, and in a simpler model that seems to capture some of the essentials in a limit, Hartnoll et.al.,[13] have studied a neutral black hole with a charged scalar and Maxwell sector that do not back react on the geometry. The latter authors have explored (with the aid of that simplifying limit) some of the phenomenology of the condensate as a function of temperature and shown that it maps rather well (where the limit can be trusted) to familiar features of superconductivity in the dual 2+1 dimensional theory.

Emboldened by these studies, we explored the case of adding an external magnetic field to the system, to see how the condensate behaves.² We have a fully back-reacted electrically and magnetically charged Reissner-Nordström black hole, and a charged scalar whose back-reaction we neglect in our computations. Since the scalar does not back-react, we cannot hope to see all of the signature physics of a superconductor in the presence of magnetism, as the superconductor is not able to repel the background magnetic field. Instead, we find that the condensate generically adjusts itself so as to fill only a strip of finite width in the plane, thereby reducing the total magnetic field that threads it. Remarkably, we can solve exactly for the profile that it adopts, and we find that, as the magnetic field approaches infinity, the condensate shrinks to zero size.

2. The background

We begin by introducing a charged, complex scalar field into the four dimensional Einstein-Maxwell action with a negative cosmological constant:³

$$S = \frac{1}{2\kappa_4^2} \int d^4x \sqrt{-G} \left\{ R + \frac{6}{L^2} + L^2 \left(-\frac{1}{4} F^2 - |\partial\Psi - igA\Psi|^2 - V(|\Psi|) \right) \right\} . \quad (2.1)$$

This action contains a term proportional to $A_\mu A^\mu \bar{\Psi}\Psi$. This term contributes negatively to the effective mass of the charged scalar since the charged black hole will source A_t . It is exactly this term that allows (but does not guarantee) a non-trivial vev for the scalar field to form. When a vev is formed, by the usual Higgs-Anderson mechanism, the gauge field develops a mass term proportional to $A_\mu A^\mu \langle \bar{\Psi}\Psi \rangle$. In the limit where the scalar field Ψ does not back-react on the geometry, the solution for the background geometry we take is that of the dyonic black hole [15]:

$$ds^2 = \frac{L^2 \alpha^2}{z^2} (-f(z) dt^2 + dx^2 + dy^2) + \frac{L^2}{z^2} \frac{dz^2}{f(z)},$$

$$F = 2h\alpha^2 dx \wedge dy + 2q\alpha dz \wedge dt,$$

$$f(z) = 1 + (h^2 + q^2) z^4 - (1 + h^2 + q^2) z^3 = (1 - z) (z^2 + z + 1 - (h^2 + q^2) z^3) . \quad (2.2)$$

In the coordinate system used in equation (2.2), z is a dimensionless radial coordinate scaled so that the event horizon of the black hole is located at $z_h = 1$ and the AdS boundary

²As we were preparing this manuscript, a paper on the same subject (ref. [14]) appeared on the ArXiv.

³We are using the mostly positive signature convention.

is at $z \rightarrow 0$. The parameters α, h , and q are related to the mass, magnetic charge, and electric charge of the black hole respectively, but only α is dimensionful, with dimension of inverse length. These quantities are in turn related to the temperature, external magnetic field, and charge density of the charged adjoint matter in the dual field theory. The only other dimensionful parameter in the solution is L , related to the AdS radius. The Hawking temperature is given by the usual Gibbons-Hawking calculus [16]:

$$T = \frac{1}{\beta} = \frac{\alpha}{4\pi} (3 - h^2 - q^2) . \tag{2.3}$$

Note that in order for the temperature to remain positive, $(h^2 + q^2) \leq 3$. Saturating this inequality corresponds to the extremal, zero-temperature case. In order to determine the effect of the magnetic and electric charges of the black hole, we choose a particular form for the gauge field A , such that $F = dA$:

$$A = 2h\alpha^2 xdy + 2q\alpha (z - 1) dt . \tag{2.4}$$

We have explicitly added the pure gauge term $-2q\alpha dt$ in order to have A regular at the event horizon [17]. The A_y term provides a constant magnetic field $B = F_{xy}$, and this is interpreted as corresponding to an external magnetic field in the (2+1)-dimensional system [18]:

$$B = 2h\alpha^2 . \tag{2.5}$$

The A_t term has two terms, a constant term and a term that goes to zero at the boundary. The constant term is interpreted as the chemical potential (for an analogue of R-charge; see e.g., refs [19, 20]), and the second term can be related to the conjugate dual charge density of the theory *via*:

$$\rho = \frac{1}{\mathcal{V}\beta} \frac{\delta S_{\text{on-shell}}}{\delta A_t(z=0)} = -\frac{L^2}{\kappa_4^2} q\alpha^2 , \tag{2.6}$$

where \mathcal{V} is the volume of the two-dimensional spatial part of the field theory.

3. The scalar field

3.1 Review

Let us review the results of ref. [13] to better clarify the relationship to our present work. In that paper, the background is neutral, so both the electric and magnetic charge of the dyonic black hole have been set to zero. Instead, the Maxwell-scalar sector is decoupled from the gravity sector by sending the coupling $g \rightarrow \infty$. In order to see this, we must first rescale $A_\mu \rightarrow A_\mu/g$ and $\Psi \rightarrow \Psi/g$. The Maxwell-scalar sector then has an overall g^{-2} , which when sent to infinity, decouples it from the gravity sector. In this analysis, the potential is taken to be:

$$V(|\Psi|) = -2\bar{\Psi}\Psi/L^2 . \tag{3.1}$$

Therefore, one can now study the Maxwell-scalar theory in the black hole background with Lagrangian:

$$\mathcal{L} = -\frac{1}{4}F^2 - |\partial\Psi - iA\Psi|^2 + 2\bar{\Psi}\Psi/L^2 \tag{3.2}$$




Figure 1: The vev of the $\Delta = 2$ operator as a function of the temperature for the isolated Maxwell-scalar sector studied in ref. [13]. T_c is proportional to the square root of the charge density. Note that we use a different normalization, which accounts for the difference of a factor of $\sqrt{2}$ with ref. [13].

The equation of motion for the fields Ψ and A_μ are:

$$\frac{1}{\sqrt{-G}} \partial_\mu \left(\sqrt{-G} G^{\mu\nu} (\partial_\nu \Psi - i A_\nu \Psi) \right) + \frac{2}{L^2} \Psi - i G^{\mu\nu} A_\mu (\partial_\nu \Psi - i A_\nu \Psi) = 0, \quad (3.3)$$

$$\frac{1}{\sqrt{-G}} \partial_\nu \left(\sqrt{-G} G^{\nu\lambda} G^{\mu\sigma} F_{\lambda\sigma} \right) - G^{\mu\nu} (i (\bar{\Psi} \partial_\nu \Psi - \partial_\nu \bar{\Psi} \Psi) + 2 A_\nu \bar{\Psi} \Psi) = 0, \quad (3.4)$$

and that of $\bar{\Psi}$ is simply the complex conjugate of equation (3.3). We take the ansatz:

$$\Psi \equiv \Psi(z) = \tilde{\Psi}(z)/L, \quad A_t \equiv A_t(z) = \alpha \tilde{A}_t(z), \quad (3.5)$$

where $\tilde{\Psi}$ and \tilde{A}_t are dimensionless fields. It is then consistent to take the phase of Ψ to be constant. All other fields are set to zero. Under this ansatz, the equations of motion simplify to:

$$\partial_z^2 \tilde{\Psi} + \left(\frac{f'}{f} - \frac{2}{z} \right) \partial_z \tilde{\Psi} + \frac{1}{f^2} \tilde{\Psi} \tilde{A}_t^2 + \frac{2}{z^2 f} \tilde{\Psi} = 0, \quad \partial_z^2 \tilde{A}_t - \frac{2}{z^2 f} \tilde{\Psi}^2 \tilde{A}_t = 0. \quad (3.6)$$

Without presenting the details of the analysis (see ref. [13]), we show in figure 1 the results of the variation of an order parameter as the temperature changes. The onset of superconductivity occurs for $T < T_c$. The critical temperature T_c is proportional to the square root of the charge density.

3.2 Perturbative limit

We now consider the scalar field as a perturbation about the dyonic black hole background. In this analysis, the Maxwell-scalar sector is not isolated from the gravity sector, since the Maxwell field has back-reacted on the gravitational background. We use the same potential considered in equation (3.1), which corresponds to choosing $m^2 L^2 = -2$ for the scalar field.

Before proceeding with our analysis, we would like to emphasize the relationship between this work and that of ref. [13], which we reviewed in the previous section. We work in the limit where the scalar does not backreact on the Maxwell fields, which should correspond approximately to taking $A_t \gg \Phi$ in ref. [13]. From equation (3.5), we see that this limit can be accomplished by taking $\alpha L \rightarrow \infty$. In this limit, the charge density diverges, and hence the limit corresponds to taking $T/T_c \rightarrow 0$, i.e. the left most end of the curve in figure 1. This argument is further established by the fact that, in this regime, A_t in the coupled equations studied in ref. [13] and reviewed earlier behaves almost linearly. In the dyonic black hole background, A_t is linear. This suggests that the analysis we propose captures the physics at $T/T_c \rightarrow 0$ in ref. [13]. Therefore, for the physics we uncover, we are well below the critical temperature.

The equation of motion for the scalar field is given by:

$$\frac{1}{\sqrt{-G}} \partial_\mu \left(\sqrt{-G} G^{\mu\nu} (\partial_\nu \Psi - ig A_\nu \Psi) \right) + \frac{2}{L^2} \Psi - ig G^{\mu\nu} A_\mu (\partial_\nu \Psi - ig A_\nu \Psi) = 0 . \quad (3.7)$$

The equation of motion for $\bar{\Psi}$ is simply the complex conjugate of equation (3.7). Using the fact that we only have A_t and A_y , and the only dependence is on the coordinates x and z , we consider an ansatz of the form $\Psi \equiv \Psi(x, z)$. The equation of motion simplifies to:

$$\frac{1}{\sqrt{-G}} \partial_z \left(\sqrt{-G} G^{zz} \partial_z \Psi \right) + G^{xx} \partial_x^2 \Psi + \frac{2}{L^2} \Psi - G^{yy} g^2 A_y^2(x) \Psi - G^{tt} g^2 A_t^2(z) \Psi = 0 . \quad (3.8)$$

This equation and its complex conjugate are purely real. Therefore, the equations of motion imply that the phase of Ψ is constant, and so without loss of generality, we take Ψ to be real. We assume a separable form for Ψ :

$$\Psi = X(x) Z(z) , \quad (3.9)$$

which further simplifies the equation of motion to:

$$\begin{aligned} \frac{1}{\sqrt{-G}} \partial_z \left(\sqrt{-G} G^{zz} Z'(z) \right) + \frac{2}{L^2} Z(z) - G^{tt} g^2 A_t^2(z) Z(z) \\ + \frac{Z(z) G^{xx}}{X} (X''(x) - g^2 A_y^2(x) X(x)) = 0 , \end{aligned} \quad (3.10)$$

where we have used the fact that $G^{xx} = G^{yy}$. In order for this equation to be consistent, we must have that:

$$X''(x) - g^2 A_y^2(x) X(x) = -k^2 X(x) , \quad (3.11)$$

where k^2 is a constant. By changing to a dimensionless variable $\tilde{x} = \sqrt{4gh\alpha^2} x$ and setting $X(x) = \tilde{X}(\tilde{x})$, equation (3.11) can be brought to the form:

$$\tilde{X}''(\tilde{x}) - \frac{\tilde{x}^2}{4} \tilde{X}(\tilde{x}) = -\frac{\tilde{k}^2}{2} \tilde{X}(\tilde{x}) , \quad (3.12)$$

where $\tilde{k}^2 = k^2/2gh\alpha^2$. Generically, the solutions to this equation can be written in terms of confluent hypergeometric functions, but for the case when \tilde{k}^2 is an odd integer, the solutions can be written in terms of Hermite functions H_n of order $n = (\tilde{k}^2 - 1)/2$. We




Figure 2: Some Hermite functions H_n of order $n = (\tilde{k}^2 - 1)/2$, for $n = 1, 3, 5$ and 101 .

restrict ourselves to this case since the Hermite functions decay exponentially for large x , which seems to be the natural physical choice. Henceforth in this paper, \tilde{k}^2 is taken to be an odd integer, and we display some examples of the functions in figure 2. However, in our present setup, we have assumed that the phase of the scalar field remains constant. Since only $\tilde{k}^2 = 1$ corresponds to a configuration that preserves its sign, it is the only physical solution for our ansatz. As a result, our equation is exactly Schrödinger's equation for a simple harmonic oscillator! Interestingly, the system confines the x -extent of the condensate in a potential that is exactly quadratic. With the constant B -field passing through the (x, y) plane, we can compute the flux of B through a region as:

$$\Phi = B\Delta x\Delta y, \tag{3.13}$$

Using the dimensionless variables we introduced earlier, we have:

$$\Phi = 2h\alpha^2 \frac{\Delta\tilde{x}\Delta\tilde{y}}{4\alpha^2 gh} = \frac{\Delta\tilde{x}\Delta\tilde{y}}{2g}. \tag{3.14}$$

$\Delta\tilde{x}$ only depends on \tilde{k}^2 since it is entirely determined by the behavior of the Hermite function. To estimate $\Delta\tilde{x}$, we use that:

$$\langle\tilde{x}^2\rangle = \int_{-\infty}^{\infty} d\tilde{x} |H_n(\tilde{x})|^2 \tilde{x}^2 = n + \frac{1}{2} = \frac{\tilde{k}^2}{2}, \tag{3.15}$$

and $\langle x \rangle = 0$, then our flux becomes:

$$\frac{\Phi}{\ell} = \frac{1}{2g} \sqrt{\frac{\tilde{k}^2}{2}}. \tag{3.16}$$

where ℓ is the extent in the y -direction, and since it is infinite there, we consider the flux per unit y -length.

Substituting our result of equation (3.11) for the x -dependence back into equation (3.10), the equation of motion for $Z(z)$ becomes:

$$\frac{1}{\sqrt{-G}} \partial_z \left(\sqrt{-G} G^{zz} Z'(z) \right) + \frac{2}{L^2} Z(z) - G^{tt} g^2 A_t^2(z) Z(z) - 2gh\alpha^2 \tilde{k}^2 Z(z) G^{xx} = 0, \quad (3.17)$$

Substituting in the functions, we get:

$$Z''(z) + \left(\frac{f'(z)}{f(z)} - \frac{2}{z} \right) Z'(z) + \left(\frac{4g^2 q^2}{f(z)^2} (z-1)^2 + \frac{2}{z^2 f(z)} - \frac{2gh\tilde{k}^2}{f(z)} \right) Z(z) = 0. \quad (3.18)$$

It is interesting to note that all α and L dependence in the equation have cancelled. This would seem to indicate that the scalar field's behavior does not depend on α and L , but it is important to remember that we are working in the perturbative limit where Ψ is supposed to be small. Earlier, we stated that to ensure the perturbative limit is consistent, we must have the quantity αL to be large. Since the only dimensionful parameters are α and L , we take:

$$\Psi = \frac{1}{\alpha L^2} \tilde{\Psi} = \frac{1}{\alpha L^2} X(x) \tilde{Z}(z), \quad (3.19)$$

which ensures that the scalar field is always perturbative, and we can now work in terms of the dimensionless function \tilde{Z} .

Near the AdS boundary, equation (3.18) becomes:

$$\tilde{Z}''(z) - \frac{2}{z} \tilde{Z}'(z) + \frac{2}{z^2} \tilde{Z}(z) = 0, \quad (3.20)$$

which has the solution:

$$\lim_{z \rightarrow 0} \tilde{Z}(z) = \Psi_1 z + \Psi_2 z^2, \quad (3.21)$$

where Ψ_1 and Ψ_2 are dimensionless constants. Both of these solutions are normalizable, so one is not the source of the other. Having chosen $m^2 L^2 = -2$ by our choice of potential (see equation (3.1)), there is not a unique boundary condition at the AdS boundary [21]. Ψ_1 is proportional to the vev of the $\Delta = 1$ operator ($\langle \mathcal{O}_1 \rangle$) and Ψ_2 is proportional to the vev of the $\Delta = 2$ operator ($\langle \mathcal{O}_2 \rangle$) in the boundary field theory, and only one of these is turned on by the boundary condition. This forces us to pick as one of our boundary conditions:

$$\Psi_1 = 0 \text{ or } \Psi_2 = 0. \quad (3.22)$$

We choose to only work with the $\Delta = 2$, since we find the same general qualitative behavior for both operators. The exact relationship between the vev and the asymptotic value of \tilde{Z} can be calculated using the holographic dictionary (details are shown in appendix A):

$$\langle \mathcal{O}_2 \rangle = \frac{\delta S_{\text{on-shell}}}{\delta \Psi(z=0)} = \frac{L^2}{2\kappa_4^2} \alpha^2 \Psi_2. \quad (3.23)$$

We can also study the behavior of the solution near the event horizon. We find that there are three distinct possible cases. First, if $gh\tilde{k}^2 < 1$, then the equation of motion becomes:

$$\tilde{Z}''(z) - \frac{1}{1-z} \tilde{Z}'(z) + \frac{a^2}{1-z} \tilde{Z}(z) = 0, \quad (3.24)$$

where

$$a^2 = 2 \frac{1 - gh\tilde{k}^2}{3 - h^2 - q^2} > 0,$$

which has Bessel functions as solutions:

$$\begin{aligned} \lim_{z \rightarrow 1} \tilde{Z}(z) &= \psi_1 J_0 \left(2a\sqrt{(1-z)} \right) + \psi_2 Y_0 \left(2a\sqrt{(1-z)} \right) \\ &\approx \psi_1 + \psi_2 \left(\frac{2}{\pi} \gamma + \frac{1}{\pi} \log(a(1-z)) \right), \end{aligned} \quad (3.25)$$

where γ is the Euler-Mascheroni constant. Since we want the field to be finite at the event horizon, we choose our other boundary condition to be $\psi_2 = 0$. For $gh\tilde{k}^2 = 1$, the equation of motion becomes

$$\tilde{Z}''(z) - \frac{1}{1-z} \tilde{Z}'(z) + b^2 \tilde{Z}(z) = 0, \quad (3.26)$$

where

$$b^2 = \frac{4g^2q^2}{(3-h^2-q^2)^2} + \frac{4}{3-h^2-q^2} > 0,$$

which has as solutions:

$$\begin{aligned} \lim_{z \rightarrow 1} \tilde{Z}(z) &= \psi_1 J_0(b^2(1-z)) + \psi_2 Y_0(b^2(1-z)) \\ &\approx \psi_1 + \psi_2 \left(\frac{2}{\pi} \gamma + \frac{2}{\pi} \log(b^2(1-z)) \right). \end{aligned} \quad (3.27)$$

Since we want the field to be finite at the event horizon, we would choose the same boundary condition as before, i.e. $\psi_2 = 0$. Finally, if $gh\tilde{k}^2 > 1$, we have:

$$\tilde{Z}''(z) - \frac{1}{1-z} \tilde{Z}'(zq) - \frac{a^2}{1-z} \tilde{Z}(z) = 0, \quad (3.28)$$

where $a^2 = 2 \frac{1-gh\tilde{k}^2}{3-h^2-q^2} > 0$, which has as solutions:

$$\begin{aligned} \lim_{z \rightarrow 1} \tilde{Z}(z) &= \psi_1 I_0 \left(2a\sqrt{(1-z)} \right) + \psi_2 K_0 \left(2a\sqrt{(1-zq)} \right) \\ &\approx \psi_1 + \psi_2 (-\gamma - \log(a(1-z))) . \end{aligned} \quad (3.29)$$

Again, if we want the field to be finite at the event horizon, we would choose the same boundary condition as before, i.e., $\psi_2 = 0$.

4. Numerical method and results

There are several parameters that we can vary in this problem, g , \tilde{k}^2 , q , h . For the following analysis, we take $g = 1$ for simplicity, but we can expect similar behavior for other values of g . In order to solve equation (3.18), we use a shooting method. We impose the following initial conditions at the event horizon:

$$\tilde{Z}(1) = 1, \quad \tilde{Z}'(1) = \frac{2 - 2gh\tilde{k}^2}{3 - h^2 - q^2}. \quad (4.1)$$




Figure 3: Two possible solutions for the scalar field with \mathcal{O}_2 turned on. The parameters used are $\tilde{k}^2 = 1$ and $h = 0.1$.

We find that these initial conditions do not necessarily satisfy equation (3.22). For a given \tilde{k}^2 , only for certain values of h and q do we get to satisfy the appropriate boundary conditions. Therefore, we fix the value of h , and then scan through possible values of q until the appropriate boundary condition at $z = 0$ is satisfied, exhibiting our condensate.

We show in figure 3 two allowed solutions for the scalar field for a fixed \tilde{k}^2 and h but with different q values. Ref. [11] argues that only the zero-node solution matters to the phase structure, so we consider only these solutions in our subsequent analysis.⁴ It is important to emphasize that changing the value of the scalar at the event horizon from the value given in equation (4.1) does not change the required value of h and q for the scalar to condense, but it does change the value of the vev. Therefore, at particular values of h and q that allow for a condensate to form, there is a whole range of allowed vev values for the operator depending on its value at the event horizon. For convenience, we define a temperature \tilde{T} (with its corresponding \tilde{q} for which the scalar condenses) to be the temperature at zero magnetic field ($h = 0$) at which we find a solution. This is given by (cf equation (2.3)):

$$\tilde{T} = \alpha \frac{3 - \tilde{q}^2}{4\pi}. \tag{4.2}$$

Using this definition, we can define several dimensionless quantities of interest:

$$\frac{\kappa_4}{L} \frac{\sqrt{\langle \mathcal{O}_2 \rangle}}{\tilde{T}} = \frac{4\pi}{3 - \tilde{q}^2} \sqrt{\frac{\Psi_2}{2}}, \quad \frac{B}{\tilde{T}^2} = \left(\frac{4\pi}{3 - \tilde{q}^2} \right)^2 2h, \quad -\frac{\kappa_4^2}{L^2} \frac{\rho}{\tilde{T}^2} = \left(\frac{4\pi}{3 - \tilde{q}^2} \right)^2 q. \tag{4.3}$$

We present the results of our numerical condensate search in terms of these quantities in figure 4.

5. Discussion

As previously alluded to, since we are working in a limit where the scalar is not back-reacting on the Maxwell field, we cannot (as in ref. [13]) track the dependence of the vev

⁴Indeed, we find that it is only for these zero-node solutions that persist for low enough temperature.




Figure 4: The graphs depict the following: the allowed B and T values for \mathcal{O}_2 to condense, the v_{ev} for \mathcal{O}_2 , and the allowed ρ and T values for \mathcal{O}_2 to condense.

on temperature all the way to the transition temperature⁵ T_c . We instead simply exhibit the values of the ratios of temperature, charge density, and magnetic field to \tilde{T} that allow a condensate to form in our study. This is enough to allow us to study the condensate’s spatial behavior, and we are firmly below T_c in all that we do..

There are several interesting features, as displayed in figure 4. Note that in ref. [13], the scalar condenses at a particular value of T/T_c , with $T_c \propto \rho^{1/2}$. Similarly, the scalar in our setup condenses at a particular value of ρ/\tilde{T}^2 and B/\tilde{T}^2 . There exists a minimum value of ρ/\tilde{T}^2 for the scalar to condense. Beyond these values, condensation occurs but is not believed to be stable [11], and the only allowed solution is the trivial one (see the previous section for more details).

At zero magnetic field, the condensate fills the plane and requires the lowest ratio of $\tilde{\rho}/\tilde{T}^2$. To see that it fills the plane, we recall that the profile in the x -direction for $\tilde{k}^2 = 1$ is given by:

$$X(x) = e^{-\frac{\tilde{x}^2}{4}} = e^{-gh\alpha^2 x^2} . \tag{5.1}$$

Therefore, in the limit of $h \rightarrow 0$, we have $X(x) \rightarrow 1$, i.e. there is no x -dependence. As the magnetic field is turned on and increases (reading the first and the other two plots

⁵Note that \tilde{T} is *not* the transition temperature, but merely a normalization set by the solution at $h = 0$.

from right to left as T/\tilde{T} decreases), the value of ρ/\tilde{T}^2 required for the scalar to condense steadily increases and the condensate has a finite thickness along the x -direction. At around $T/\tilde{T} = 0.25$, the ratio ρ/\tilde{T}^2 needed drops rapidly. The condensate also drops rapidly around the same interval, which suggests that the magnetic field might be overcoming the forces keeping the superconductor together. However, we find that even at $T/\tilde{T} = 0$, the condensate persists with a standard deviation of $1/\sqrt{6g\alpha^2}$.

This result is of particular interest because, in the limit of large magnetic field, i.e. $\alpha \rightarrow \infty$, the standard deviation approaches zero. Therefore, the magnetic field, as it grows, shrinks the condensate away completely. This is reminiscent of the Meissner effect, where the magnetic field expels the condensate. The condensate itself cannot expel the magnetic field, as is usually the case, since the scalar cannot back-react on the background magnetic field.

It is interesting to speculate on what higher \tilde{k}^2 values could mean if for example the ansatz for the scalar field is modified to make them physical. As an approximation, we can naively proceed with our setup with higher values of \tilde{k}^2 . We find that as \tilde{k}^2 increases and the x -width of the condensate expands (see figure 2), the corresponding B -field associated with it is smaller. It is interesting to follow this to large \tilde{k}^2 , the ‘‘classical’’ limit of the quantum harmonic oscillator controlling the x -profile: The $\tilde{k}^2 \rightarrow \infty$ limit has the condensate filling the plane while the magnetic field $B \rightarrow 0$. Pleasingly, this is consistent with the limit of small and large magnetic field we discussed above.⁶

Acknowledgments

TA thanks L’Institut de Physique Th eorique (IPhT), Saclay, France for hospitality. He would also like to thank Nikolay Bobev and Hubert Saleur for discussions. This research was supported by the U.S. Department of Energy.

A. Holographic dictionary

The solution of the equation of motion admits two normalizable solutions at the AdS boundary, which we reproduce here:

$$\lim_{z \rightarrow 0} \tilde{Z}(z) = \Psi_1 z + \Psi_2 z^2,$$

To proceed with calculating the vev of the $\Delta = 2$ operator \mathcal{O}_2 using the holographic dictionary, the procedure is to assume that Ψ_1 is the source of the operator [21]. To proceed, we first write the asymptotic solution of the full scalar field as:

$$\lim_{z \rightarrow 0} \Psi(x, z) = e^{i\varphi} (\phi_0(x)z + A(x)z^2), \tag{A.1}$$

⁶In making sense of the B -field within the condensed region, where we expect it would be forced to zero in the fully back-reacting system, it is tempting to interpret the x profile of the scalar quite literally and take B times the average value within the region. This would mostly then integrate to zero in the interior of the sample, leaving only some non-zero contribution at the edges when $n = (\tilde{k}^2 - 1)/2$ is even, and canceling exactly when n is odd, but this is possibly too naive.

where we have explicitly shown the constant phase of the scalar field. Next, we calculate the variation of the on-shell action:

$$\begin{aligned} \delta S_{\text{on-shell}} &= -\frac{L^2}{2\kappa_4^2} \int d^3x \sqrt{-G} G^{zz} \partial_z \Psi(x, z) \delta \bar{\Psi}(x, z) \Big|_{z=0}^{z=1} \\ &= \lim_{z \rightarrow 0} \frac{L^2}{2\kappa_4^2} \int d^3x \frac{L^2 \alpha^3}{z^2} (z \phi_0(x) \delta \phi_0(x) + 2z^2 A(x) \delta \phi_0(x) + z^2 \phi_0(x) \delta A(x) + O(z^3)) \end{aligned}$$

At this point, we may worry about the divergence produced by the first term in this expression, but we have not included the following counterterm in the expression of our action:

$$S_{\text{counter}} = -\frac{L^2}{4\kappa_4^2} \sqrt{-\gamma} \frac{1}{L} \int d^3x \bar{\Psi}(x, z=0) \Psi(x, z=0), \quad (\text{A.2})$$

where γ is not the Euler-Mascheroni constant. Varying this counterterm and including it in the on-shell action gives as a final result:

$$\delta S_{\text{on-shell}} = \frac{L^2}{2\kappa_4^2} \int d^3x L^2 \alpha^3 A(x) \delta \phi_0(x) \quad (\text{A.3})$$

Therefore, we have our final result:

$$\langle \mathcal{O}_2(x) \rangle = \frac{1}{d\beta} \frac{\delta S_{\text{on-shell}}}{\delta \phi_0(x)} = \frac{L^2}{2\kappa_4^2} L^2 \alpha^3 A(x) = \frac{L^2}{2\kappa_4^2} \alpha^2 \Psi_2 X(x), \quad (\text{A.4})$$

where d represents the fact that we want to study the operator in terms of unit length in the y -direction. If we wish to only consider the overall scale of the operator (i.e., dropping the x dependence), we can simply write:

$$\langle \mathcal{O}_2 \rangle = \frac{L^2}{2\kappa_4^2} \alpha^2 \Psi_2. \quad (\text{A.5})$$

References

- [1] J.M. Maldacena, *The large- N limit of superconformal field theories and supergravity*, *Adv. Theor. Math. Phys.* **2** (1998) 231 [*Int. J. Theor. Phys.* **38** (1999) 1113] [[hep-th/9711200](#)].
- [2] S.S. Gubser, I.R. Klebanov and A.M. Polyakov, *Gauge theory correlators from non-critical string theory*, *Phys. Lett.* **B 428** (1998) 105 [[hep-th/9802109](#)].
- [3] E. Witten, *Anti-de Sitter space and holography*, *Adv. Theor. Math. Phys.* **2** (1998) 253 [[hep-th/9802150](#)].
- [4] O. Aharony, S.S. Gubser, J.M. Maldacena, H. Ooguri and Y. Oz, *Large- N field theories, string theory and gravity*, *Phys. Rept.* **323** (2000) 183 [[hep-th/9905111](#)].
- [5] R.M. Wald, *Black hole in a uniform magnetic field*, *Phys. Rev.* **D 10** (1974) 1680.
- [6] J. Bicak and L. Dvorak, *Stationary electromagnetic fields around black holes. 3. General solutions and the fields of current loops near the Reissner-Nordstrom black hole*, *Phys. Rev.* **D 22** (1980) 2933.

- [7] A. Chamblin, R. Emparan and G.W. Gibbons, *Superconducting p-branes and extremal black holes*, *Phys. Rev. D* **58** (1998) 084009 [[hep-th/9806017](#)].
- [8] R. Ruffini and J.A. Wheeler, *Introducing the black hole*, *Phys. Today* **24** (1971) 30.
- [9] J.D. Bekenstein, *Black hole hair: twenty-five years after*, [gr-qc/9605059](#).
- [10] S.S. Gubser, *Phase transitions near black hole horizons*, *Class. and Quant. Grav.* **22** (2005) 5121 [[hep-th/0505189](#)].
- [11] S.S. Gubser, *Breaking an abelian gauge symmetry near a black hole horizon*, [arXiv:0801.2977](#).
- [12] S.S. Gubser, *Colorful horizons with charge in anti-de Sitter space*, [arXiv:0803.3483](#).
- [13] S.A. Hartnoll, C.P. Herzog and G.T. Horowitz, *Building a holographic superconductor*, *Phys. Rev. Lett.* **101** (2008) 031601 [[arXiv:0803.3295](#)].
- [14] E. Nakano and W.-Y. Wen, *Critical magnetic field in a holographic superconductor*, *Phys. Rev. D* **78** (2008) 046004 [[arXiv:0804.3180](#)].
- [15] L.J. Romans, *Supersymmetric, cold and lukewarm black holes in cosmological Einstein-Maxwell theory*, *Nucl. Phys. B* **383** (1992) 395 [[hep-th/9203018](#)].
- [16] G.W. Gibbons and S.W. Hawking, *Classification of gravitational instanton symmetries*, *Commun. Math. Phys.* **66** (1979) 291.
- [17] S. Kobayashi, D. Mateos, S. Matsuura, R.C. Myers and R.M. Thomson, *Holographic phase transitions at finite baryon density*, *JHEP* **02** (2007) 016 [[hep-th/0611099](#)].
- [18] S.A. Hartnoll and P. Kovtun, *Hall conductivity from dyonic black holes*, *Phys. Rev. D* **76** (2007) 066001 [[arXiv:0704.1160](#)].
- [19] A. Chamblin, R. Emparan, C.V. Johnson and R.C. Myers, *Charged AdS black holes and catastrophic holography*, *Phys. Rev. D* **60** (1999) 064018 [[hep-th/9902170](#)].
- [20] A. Chamblin, R. Emparan, C.V. Johnson and R.C. Myers, *Holography, thermodynamics and fluctuations of charged AdS black holes*, *Phys. Rev. D* **60** (1999) 104026 [[hep-th/9904197](#)].
- [21] I.R. Klebanov and E. Witten, *AdS/CFT correspondence and symmetry breaking*, *Nucl. Phys. B* **556** (1999) 89 [[hep-th/9905104](#)].